

Evidence against or for topological defects in the BOOMERanG data ?

F. R. Bouchet¹, P. Peter¹, A. Riazuelo², M. Sakellariadou^{1,3}

¹*Institut d'Astrophysique de Paris, 98bis boulevard Arago, 75014 Paris, France*

²*Université de Genève, Service de Physique Théorique, 24 Quai Ernest Ansermet, 1211 Genève 4, Switzerland*

³*Section of Astrophysics, Astronomy and Mechanics, University of Athens, Panepistimiopolis, 15784 Zografos, Hellas*

The recently released BOOMERanG data was taken as “contradicting topological defect predictions”. We show that such a statement is partly misleading. Indeed, the presence of a series of acoustic peaks is perfectly compatible with a non-negligible topological defects contribution. In such a mixed perturbation model (inflation and topological defects) for the source of primordial fluctuations, the natural prediction is a slightly lower amplitude for the Doppler peaks, a feature shared by many other purely inflationary models. Thus, for the moment, it seems difficult to rule out these models with the current data.

For almost two decades, two families of models have been considered challengers for describing, within the framework of gravitational instability, the formation of large-scale structure in the universe. Initial density perturbations can either be induced by quantum fluctuations of a scalar field during inflation [1], or they may be triggered by a class of topological defects [2]. The inflationary paradigm was proposed in order to explain the shortcomings of the standard Big Bang model, namely the horizon and flatness problems. Topological defects, on the other hand, can be formed during symmetry-breaking phase transitions in the early universe [3]. Moreover they are naturally predicted by many realistic models of particle physics aiming at describing interactions at energies much higher than what is currently reachable in accelerators [2].

The Cosmic Microwave Background (CMB) anisotropies provide a powerful tool to discriminate among inflation and topological defects. CMB anisotropies are conveniently characterized by their angular power spectrum C_ℓ , the average value of the square of the coefficients, $|a_{\ell m}|^2$, of a spherical harmonic decomposition of the measured CMB pattern. This spectrum fully characterizes a normal distributed field, and $\ell(\ell+1)C_\ell$ is the logarithmic contribution to the variance of multipoles ℓ .

On large angular scales ($\ell \lesssim 50$), both families of models lead to approximately scale-invariant spectra, with however a different prediction regarding the statistics of the induced perturbations. Provided the quantum fields are initially placed in the vacuum, inflation predicts generically Gaussian fluctuations, whereas in the case of topological defects models, the induced perturbations are clearly non-Gaussian, at least at sufficiently high angular resolution. This is an interesting fingerprint, even though difficult to test through the data. In the context of inflation, non-Gaussianity can however also be present, as for example in the case of stochastic inflation [4], or in a class of inflationary models involving two scalar fields leading to non-Gaussian isothermal fluctuations with a blue spectrum [5]. In addition, allowing non-vacuum initial states for the cosmological perturbations of quantum-mechanical origin, one generically obtains a non-Gaussian spectrum [6]. Finally, in a cosmological model where perturbations are induced by

inflation with a non-negligible topological defects contribution, one again expects deviation from Gaussianity.

Both inflation and defect models predict a spectrum of temperature anisotropies at the required 10^{-5} level [7] on large angular scales. However, on intermediate and small angular scales, the predictions of inflation are quite different than those of topological defects, due to the different nature of the induced perturbations. More precisely, the inflationary fluctuations are coherent, in the sense that the perturbations are initially at the same phase and subsequently evolve linearly and independently of each other. The subsequent progressive phase shift between different modes thus produces the so-called acoustic (or Doppler) peak structure. On the other hand, in topological defect models, fluctuations are constantly induced by the sources (defects). Since topological defects evolve in a non-linear manner, and since the random initial conditions of the source term in the perturbation equations of a given scale leaks into other scales, perfect coherence is destroyed [8,9].

In the inflationary case, coherent adiabatic fluctuations lead to a rather high first acoustic peak, around $\ell \sim 200$ (for $\Omega \sim 1$) followed by a set of oscillatory peaks for larger ℓ . As a matter of illustration, in the Standard Cold Dark Matter (SCDM) model with $n_s = 1$, $h = 0.5$, $\Omega_0 = 1$, $\Omega_b = 0.05$, and vanishing cosmological constant, the second peak is expected to be three times higher than the low ℓ (Sachs-Wolfe) plateau.

On the other hand, in topological defect models, incoherent fluctuations lead to a single bump at smaller angular scales (larger ℓ). For instance, global $O(4)$ textures predict a peak whose position is shifted to $\ell \simeq 350$ having an amplitude which is ~ 1.5 times higher than the Sachs-Wolfe plateau [10]. Global $O(N)$ textures in the large N limit lead to a rather flat spectrum, with a slow decay after $\ell \sim 100$ [11]. Roughly similar results are obtained with other global $O(N)$ defects [12,13]. Local cosmic string predictions are unfortunately not very well established in detail and range from an almost flat spectrum [14] to a single wide bump at $\ell \sim 500$ [15] with extremely rapidly decaying tail. It seems that the microphysics of the string network plays a crucial role in the height and in the position of the bump [16,17]. An interesting point however is that global textures (and es-

pecially global $O(N)$ defects in the large N limit, where every component of the field which produces the defects evolves almost linearly, see [11]) are in a sense “less non-linear” than local cosmic strings. This translates into the presence of a very smooth second peak in the spectrum.

Before the first Antarctic BOOMERanG [18] results, several experiments had already given some hints for a rise in the power spectrum at the degree scale ($\ell \sim 200$). Then, more recently, the TOCO [19] experiment and the technical flight over North America of BOOMERanG itself [20] detected a maximum in the spectrum around the same scale, followed by a subsequent decay, thereby ruling out pure topological defect based scenarios. But measurements at $\ell > 400$ were rather crude (e.g., [21]). The BOOMERanG experiment has spanned in a single experiment a very wide range of angular scales, from $\ell \simeq 50$ to $\ell \simeq 800$ and has reached sufficient sensitivity to map out the second and third peak region of the spectrum.

Since then, the observational situation has dramatically improved. The analysis of the BOOMERanG-LDB data set, which is exceptional in data quality, span of scales covered, and angular resolution, firmly reveals the presence of a relatively narrow peak in the power spectrum at $\ell \sim 200$ followed by an almost flat plateau, roughly at the same level as the Sachs-Wolfe part. The first peak is found to be at the position expected in the case of a flat universe.

The data at higher ℓ lead to some confusion. In the first release of the data [18], only a relatively low secondary peak could be accommodated. Even though there was no direct measurement of the height of the second peak, one could still, safely enough, remark that the relative amplitude between the two peaks was rather high. Indeed it was well above its expected value, even in the case of a high baryonic density universe which tends to enhance the contrast between odd and even peaks (see [22] for a detailed description of the influence of cosmological parameters on the CMB anisotropy spectrum). This finding has led to new directions for the explanation of the CMB anisotropies. We briefly discuss some suggestions below. Within the context of inflationary perturbations, the data required [23] a tilted model with a red spectrum (neglecting the possible contribution of tensor modes, i.e., gravitational waves, the measured spectral index would appear to be of order $n_s \simeq 0.85$) and/or a very high baryonic density, more than one standard deviation away from the nucleosynthesis constraint [24]. Such conclusions put the simplest inflationary scenario in difficulty. Other scenarios were then proposed in order to explain a relatively low second peak together with a high and narrow first one, such as, e.g., including a leptonic asymmetry [25]. Another possibility was to add power at low ℓ . There are several possibilities to do so, for example by adding a tensor contribution (which is rather flat before $\ell \simeq 100$ and rapidly decays afterwards; note that if one considers an inflationary scenario with a red spectral index, there is necessarily such a non negligible tensor contribution), or isocurvature fluctuations [26,27].

These possibilities were discussed in Ref. [28].

The more recent BOOMERanG data [40] somewhat changed this picture. A new estimation of the beam size and of the calibration uncertainties, led to a significant increase of the power spectrum in the region of the secondary peaks, as well as a confirmation of the long expected acoustic oscillations in the spectrum. Then, the estimation of the cosmological parameters from the shape of the spectrum led to values definitely more compatible with the currently ones derived by many independent astrophysical measurements. Although there are still some differences with the new MAXIMA data [41], this might be seen as the long-awaited advent of “consistent cosmology” [42]. However, in order to give firm bounds on the value of the cosmological parameters, we still must ensure that we have explored the correct region of the parameter space. Since there already exist very good fits to the data, this reduces to ask at what extent the currently favored set of cosmological parameters might be degenerate with other yet non explored but physically relevant parameters.

In this article, we illustrate a new degeneracy arising in the data, that would be due to a small but significant contribution of topological defects. These mixed perturbation models (i.e., both of inflationary type and seeded by topological defects) have already been proposed on other grounds. Indeed, the formation of topological defects provides the necessary mechanism in order to successfully exit the inflationary era [29] in a number of particle physics motivated inflationary [30–32] models. Within the context of hybrid inflation [33] in supergravity, a model leading to a scenario where CMB anisotropies may be produced by cosmic strings on a $\sim 10^{16}$ GeV mass scale, and galaxy formation may be due to inflationary perturbations, was proposed in Ref. [31] and further investigated in Refs. [34,35]. In addition, in most classes of superstring compactification involving the spontaneous breaking of a pseudo-anomalous $U(1)$ gauge symmetry [36], strings of the cosmic kind [37] are also formed [38] and turn out to be local [39]. We therefore consider a model in which a network of cosmic strings evolved [43] independently of any pre-existing fluctuation background generated by a standard cold dark matter with a non-zero cosmological constant (Λ CDM) inflationary phase. As we shall restrict our attention to the angular spectrum, we can remain in the linear regime all along our analysis, so that setting C_ℓ^I and C_ℓ^S the (COBE normalized) Legendre coefficients due to adiabatic inflation fluctuations and those stemming from the string network respectively, we have

$$C_\ell = \alpha C_\ell^I + (1 - \alpha) C_\ell^S. \quad (1)$$

In the above equation, the C_ℓ are to be compared with the observed data and the coefficient α is a free parameter giving the relative amplitude for the two contributions. In this preliminary work, we do not vary C_ℓ^S characteristics and simply use the model of Ref. [13]. Strictly speaking, the anisotropy power spectrum reported in Ref. [13]

concerns theories of global defects. However, the finding that vector and tensor modes provide substantial contribution to the large angular scale anisotropies, hold for both global defect models [12,13] and cosmic string scenarios [14]. Moreover, models of global strings as well as of local strings, are characterized by incoherent perturbations. We thus believe that the spectrum of Ref. [13] exemplifies the power spectra of both global and local cosmic strings. In addition, another advantage of this model, is that it takes into account the expansion of the universe. To obtain the power spectrum from numerical simulations with cosmic strings, one must take into account the “three-scale model” [44] of cosmic string networks, the small-scale structure (wiggleness) [17] of the strings, the microphysics of the string network [16], as well as the expansion of the universe. This is a rather difficult task and to our knowledge no currently available simulation includes all of them.




FIG. 1. $\ell(\ell+1)C_\ell$ versus ℓ for three different models. The upper dot-dashed line represents the prediction of a Λ CDM model, with cosmological parameters set as $n_s = 1$, $\Omega_\Lambda = 0.7$, $\Omega_m = 0.3$, $\Omega_b = 0.05$ and $h = 0.6$ in agreement with all other data but CMB’s. The lower dashed line is a typical string spectrum. Both of these are seen not to fit the new BOOMERanG, MAXIMA and DASI data (circles, triangles and diamonds respectively) and are normalized at the COBE scale (crosses). Combining both curves with the extra-parameter α produces the solid curve, with a χ^2 per degree of freedom slightly above unity. The string contribution turns out to be some 18% of the total. With the former BOOMERanG data which produced a much lower second peak, the string content was raised to 38% of the total.

Figure 1 shows the two uncorrelated spectra as a function of ℓ , both normalized on the COBE data, together with the weighted sum. The best fit, having $\alpha \sim 0.82$ yields a contribution in strings which is not negligible, although the inflation-produced perturbations definitely represent the dominant part for this spectrum. Also, our conclusions are not significantly affected if we remove the

new MAXIMA [41] and DASI [45] data. The fact that the string content extracted from the data is not very large is of course related to the fact that our choice of value for the cosmological parameters are very similar to the best fit found by the BOOMERanG team [40]. For example, the string content would be larger should we consider the former BOOMERanG 2000 [18] data instead. Nevertheless, it illustrates that there is some degree of degeneracy between this model with a string contribution and the model without strings even with popular cosmological parameters and “good” data. It thus seems to us necessary to add the string contribution as a new parameter to the standard parameter space. Actually, it appears quite natural to enlarge the parameter space as increasingly high quality data begin to be at our disposal. In the same vein, there seems to be a growing agreement on the fact that a tensor contribution has to be taken into account when trying to fit the data, and that the presence of non adiabatic initial conditions also deserves to be more carefully studied.

We would like to briefly comment upon the “biasing problem” in structure formation. Topological defect models have been sometimes claimed to be ruled out, since they would not lead to large enough matter density perturbations, once normalized to the COBE data on very large scales. This normalization fixes the only free parameter of a given defect model, namely the symmetry breaking scale. More precisely, on scales of $100 h^{-1}$ Mpc, which are most probably unaffected by non-linear gravitational evolution, standard topological defect models, once normalized to COBE, require a bias factor of $b_{100} \sim 5$, whereas b_{100} is most probably close to unity. In cosmic string models, the biasing problem is definitely not that severe, since it is model dependent. More precisely, the matter power spectrum is very sensitive to the assumptions made about string decay [15,16]. Clearly, the “biasing problem” may be cured within mixed perturbation scenarios, in which structure arises by the combined effect of both adiabatic density perturbations produced at the end of inflation and isocurvature fluctuations seeded by topological defects.

In conclusion, we have found that a mixture of inflation and topological defects can perfectly well accommodate the current CMB data, with standard values of the cosmological parameters. We think that this new, unexpected, degeneracy should be carefully taken into account in the future CMB data analysis. We are well aware that the analysis presented in this paper is very coarse. For instance, the actual shape of the string spectrum which we used is yet very uncertain and a more detailed study is required. In particular, it could be parameterized so that a more thorough analysis could be performed, not only on one single variable α , but over the full cosmological parameter set in order to incorporate all the effects here neglected. In addition one would have to use the full covariance matrix of the BOOMERanG band powers rather than their quoted error bars only. However, we believe that the result shown on Fig. 1 provides an

indication that the role of topological defects in structure formation should not be underestimated just on the ground that they happen to be unable to reproduce all the data if they are the sole source of primordial fluctuations.

It is a pleasure to thank B. Carter, N. Deruelle, R. Juszkiewicz, A.D. Linde and J.-P. Uzan for enlightening comments.

-
- [1] A.H. Guth, Phys. Rev. D **23**, 347 (1981); A.D. Linde, Phys. Lett. B **108**, 389 (1982); A. Albrecht, P.J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982); A.D. Linde, *Particle Physics and Inflationary Cosmology* (Harwood Academic, New York, 1990).
 - [2] T.W.B. Kibble, J. Phys. A **9**, 1387 (1976), Phys. Rep. **67**, 183 (1980); E.P.S. Shellard and A. Vilenkin, *Cosmic strings and other topological defects* (Cambridge University Press, Cambridge, England, 1994); M.B. Hindmarsh and T.W.B. Kibble, Rep. Prog. Phys. **58**, 477 (1995).
 - [3] T.W.B. Kibble, Phys. Rep. **67**, 183 (1980).
 - [4] A. Gangui, Phys. Rev. D **50**, 3684 (1994); A. Gangui, F. Lucchin, S. Matarrese and S. Mollerach, Astrophys. J. **430**, 447 (1994); A. Matacz, Phys. Rev. D **55**, 1860 (1997).
 - [5] A. Linde and V. Mukhanov, Phys. Rev. D **56**, 535 (1997).
 - [6] J. Martin, A. Riazuelo and M. Sakellariadou, Phys. Rev. D **61**, 083518 (2000).
 - [7] G.F. Smoot *et al.*, Astrophys. J. **396**, L1 (1992); C.L. Bennett *et al.*, Astrophys. J. **464**, L1 (1996); See <http://www.sns.ias.edu/~max/index.html> for an up-to-date compilation of the current data.
 - [8] J. Magueijo, A. Albrecht, P. Ferreira and D. Coulson, Phys. Rev. D **54**, 3727 (1996).
 - [9] A. Riazuelo and N. Deruelle, Annalen der Physik (Leipzig) **9**, 288 (2000).
 - [10] R. Durrer, A. Gangui and M. Sakellariadou, Phys. Rev. Lett. **76**, 579 (1996).
 - [11] R. Durrer, M. Kunz and A. Melchiorri, Phys. Rev. D **59**, 123005 (1999).
 - [12] N. Turok, U.-L. Pen and U. Seljak, Phys. Rev. D **58**, 023506 (1998).
 - [13] U.-L. Pen, U. Seljak and N. Turok, Phys. Rev. Lett. **79**, 1611 (1997).
 - [14] B. Allen, R.R. Caldwell, S. Dodelson, L. Knox, E.P.S. Shellard and A. Stebbins, Phys. Rev. Lett. **79**, 2624 (1997).
 - [15] C. Contaldi, M. Hindmarsh and J. Magueijo, Phys. Rev. Lett. **82**, 679 (1999).
 - [16] A. Riazuelo, N. Deruelle and P. Peter, Phys. Rev. D **61**, 123504 (2000).
 - [17] L. Pogosian and T. Vachaspati, Phys. Rev. D **60**, 083504 (1999).
 - [18] P. de Bernardis *et al.*, Nature (London) **404**, 955 (2000).
 - See also <http://puck.roma1.infn.it/~>.
 - [19] E. Torbet, M.J. Devlin and W.B. Dorwart *et al.*, Astrophys. J. **521**, L79 (1999); A.D. Miller, R.R. Caldwell and M.J. Devlin *et al.*, Astrophys. J. **524**, L1 (1999).
 - [20] P.D. Mauskopf *et al.*, Astrophys. J. **536**, L59 (2000).
 - [21] P.F. Scott *et al.*, Astrophys. J. **461**, L1 (1996); J. Baker *et al.*, MNRAS **308**, 1173 (1999).
 - [22] W. Hu, N. Sugiyama and J. Silk, Nature (London) **386**, 37 (1997).
 - [23] M. White, D. Scott and E. Pierpaoli, Astrophys. J. **545**, 1 (2000).
 - [24] See, e.g., K.A. Olive, Nucl. Phys. (proc. Suppl.) **B80**, 79 (2000) [proceedings of the XIXth Texas Symposium, edited by E. Aubourg *et al.* (Elsevier, Amsterdam, The Netherlands, 2000)] and references therein.
 - [25] J. Lesgourgues and M. Peloso, Phys. Rev. D **62**, 081301 (2000).
 - [26] K. Enqvist and H. Kurki-Suonio, Phys. Rev. D **61**, 043002 (2000).
 - [27] D. Langlois and A. Riazuelo, Phys. Rev. D **62**, 043504 (2000).
 - [28] M. Tegmark and M. Zaldarriaga, Phys. Rev. Lett. **85**, 2240 (2000).
 - [29] R. Jeannerot, Phys. Rev. D **56**, 6205 (1997).
 - [30] L.A. Kofman and A.D. Linde, Nucl. Phys. **B282**, 555 (1987).
 - [31] A.D. Linde and A. Riotto, Phys. Rev. D **56**, 1841 (1997).
 - [32] D.H. Lyth and A. Riotto, Phys. Rep. **314**, 1 (1999).
 - [33] A.D. Linde, Phys. Lett. B **259**, 38 (1991); Phys. Rev. D **49**, 748 (1994).
 - [34] C. Contaldi, M. Hindmarsh and J. Magueijo, Phys. Rev. Lett. **82**, 2034 (1999).
 - [35] R.A. Battye and J. Weller, Phys. Rev. D **61**, 043501 (2000); R.A. Battye, J. Magueijo and J. Weller, astro-ph/9906093 (1999).
 - [36] M. Dine, N. Seiberg and E. Witten, Nucl. Phys. **B289**, 589 (1987); J. Atick, L. Dixon and A. Sen, Nucl. Phys. **B292**, 109 (1987); M. Dine, I. Ichinose and N. Seiberg, Nucl. Phys. **B293**, 253 (1988).
 - [37] A. Casas and C. Muñoz, Phys. Lett. B **216**, 37 (1989); A. Casas, J. Moreno, C. Muñoz and M. Quiros, Nucl. Phys. **B328**, 272 (1989).
 - [38] J.A. Harvey and S.G. Naculich, Phys. Lett. B **217**, 231 (1989).
 - [39] P. Binétruy, C. Deffayet and P. Peter, Phys. Lett. B **441**, 52 (1998).
 - [40] C.B. Netterfield *et al.*, astro-ph/0104460; P. de Bernardis *et al.*, astro-ph/0105296.
 - [41] A.T. Lee *et al.*, astro-ph/0104459; R. Stompor *et al.*, astro-ph/0105062.
 - [42] X. Wang, M. Tegmark and M. Zaldarriaga, astro-ph/0105091.
 - [43] D.P. Bennett and F.R. Bouchet, Phys. Rev. D **41**, 2408 (1990); D.P. Bennett and F.R. Bouchet, Phys. Rev. Lett. **63**, 2776 (1989); F.R. Bouchet, D.P. Bennett and A. Stebbins, Nature (London) **335** (1988).
 - [44] D. Austin, E.J. Copeland and T.W.B. Kibble, Phys. Rev. D **48**, 5594 (1993).
 - [45] N.W. Halverson *et al.*, astro-ph/0104489; C. Pryke *et al.*, astro-ph/0104490.